

**ASSIGNMENT AND PRELIMINARY ANALYSIS
OF THE IR ABSORPTION SPECTRUM
OF CH₃D IN THE REGION 1900-3200 CM⁻¹**

Nikitin A.⁽¹⁾, Champion J.P.⁽²⁾, Tyuterev V.I.G.⁽³⁾
and Brown L.R.⁽⁴⁾

(1) Laboratory of Theoretical Spectroscopy, Institute of Atmospheric Optics,
Russian Academy of Sciences, 634055 Tomsk, Russia.

(2) Laboratoire de Physique URA CNRS 1796, Université de Bourgogne, 6
Boulevard Gabriel, 21000 Dijon, France.

(3) GSMA, URA CNRS D 1434, Faculté des Sciences, BP 347, 51062 Reims,
France.

(4) Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA
91109 USA.

The infrared spectra of CH₃D in the region 1900-3200 cm⁻¹ has been investigated using high-resolution Fourier spectra recorded at Kitt Peak. This region includes the nine interacting vibrational bands: $\nu_1(A_1)$, $\nu_2(A_1)$, $\nu_4(E)$, $2\nu_3(A_1)$, $2\nu_5(A_1+E)$, $2\nu_6(A_1+E)$, $\nu_3+\nu_5(E)$, $\nu_3+\nu_6(E)$ and $\nu_5+\nu_6(A_1+A_2+E)$.

Starting from assignments available in HITRAN, the analysis process could be initiated leading on turn to an extended assignment of the new spectra involving all vibrational bands. However in the present stage, due to problems encountered with two of the bands, the fit was limited to small values of J.

The investigation of such a complex polyad (so-called Nonad) required to develop an appropriate model based on the tensorial formalism allowing a coherent description of the successive polyads of the molecule. The Hamiltonian was expressed as a sum of terms relating directly to the subsequent Polyads: Ground State, Triad and Nonad. Doing so, the effects of the rovibrational interactions taking place in the Triad system (1) are consistently extrapolated into the Nonad system. Theoretical details including relationships between our parameters and classical parameters are described elsewhere (2).

Part of the research reported in this paper was performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

(1) G.Tarrago, M.Delaveau, L.Fusina and G.Guelachvili, J. Mol. Spectrosc. 126, 149 (1987).

(2) A.Nikitin, J.P. Champion, and V.I.G.Tyuterev (Poster Dijon 1995).